

Вісник першої української жінки України



# БІОЛОГІЧНИЙ ВІСНИК

МДУ ім. Б. Хмельницького

**2016. 6(3)**

Biological Bulletin of Bogdan Chmelnytsky  
National State Pedagogical University



Journal of Biology. Founded in 2011, Volynsk

## Articles

- PRELIMINARY SURVEY OF THE GROUND-DWELLING ARTHROPODS OF THE FLOOD-PLAIN MEADOWS IN THE SOUTHEAST OF POLTAVA REGION (UKRAINE) 5  
N. Zhuravel, N. Polchaninova, I. Lezhenina, O. Drovgalenko, A. Putchkov
- ОЦІНКА АДАПТИВНОЇ ЗДАТНОСТІ ТА ЕКОЛОГІЧНОЇ СТАБІЛЬНОСТІ СКОРОСТИГЛИХ ГІБРИДІВ КУКУРУДЗИ (*ZEA MAYS* L.) В КОНТРАСТНИХ УМОВАХ ВИПРОБУВАННЯ 18  
V. Yu. Cherchel, O. L. Gaidash
- MATERIALS TO THE SPIDER FAUNA (ARANEI) OF THE BYKOVA SHEYA SITE OF THE "GALICH' YA GORA" NATURE RESERVE (LIPETSK REGION, RUSSIA) 26  
N. Yu. Polchaninova
- УРОВНИ ПЛОИДНОСТІ І ОТНОСИТЕЛЬНОГО СОДЕРЖАННЯ ДНК В КУЛЬТУРЕ КЛЕТОК І ТКАНЕЙ РАСТЕНІЙ IN VITRO 33  
M. V. Skaptsov, M. A. Krasnoborodkina, M. G. Kutsev, S. V. Smirnov, A. I. Shmakov, A. V. Matsyura
- КОНСПЕКТ РОДА АСТІНОПТЕРІС ЛІНК (PTERIDACEAE) 39  
A. V. Vaganov, A. I. Shmakov
- REDESCRIPTION OF GENUS *PSYCHONOSTUA* GROTE, 1865 (INSECTA: LEPIDOPTERA, COSSIDAE, ZEUZERINAE) 46  
R. V. Yakovlev, F. C. Penco, T. J. Witt
- АГРОЕКОЛОГІЧНЕ ВИПРОБУВАННЯ СОРТІВ ЯРИХ ЗЕРНОВИХ КУЛЬТУР У ПІВНІЧНОМУ СТЕПУ УКРАЇНИ 54  
A. D. Gyrka, I. O. Kulyk, O. O. Pedash, O. O. Viniukov, V. A. Ischenko
- THE CRYPTOCEPHALINAE (COLEOPTERA: CHRYSOMELIDAE) OF THE MONGOLIAN ALTAI 61  
E. V. Guskova
- THE PTEROPHORIDAE FAUNA (LEPIDOPTERA) OF PAVLODAR REGION 73  
P. Ustjuzhanin, S. Titov, V. Kovtunovich, K. Akhmetov
- ВПЛИВ РЕГУЛЯТОРУ РОСТУ НА ФЕРМЕНТИ АНТИОКСИДАНТНОГО ЗАХИСТУ ЗА УМОВ ДІІ АНТРОПОГЕННИХ ЕМІСІЙ 80  
O. M. Vasilyuk
- ФИТОИНДИКАЦИОННОЕ ОЦЕНИВАНИЕ КАТЕНА СООБЩЕСТВ ПОЧВЕННОЙ МЕЗОФАУНЫ И ИХ ЭКОМОРФИЧЕСКАЯ ОРГАНИЗАЦИЯ 91  
A. V. Zhukov, O. N. Kunah, V. A. Novikova, D. S. Ganzha
- ГЕНЕТИЧЕСКИЙ КОНТРОЛЬ УСТОЙЧИВОСТИ К СТЕБЛЕВОЙ РЖАВЧИНЕ У ИНТРОГРЕССИВНЫХ ЛИНИЙ ТВЕРДОЙ ПШЕНИЦЫ, ПРОИЗВОДНЫХ *TRITICUM TIMORNEE* VII ЗНУК. 121  
L. P. Khlebova, N. V. Barysheva
- ЕМКОСТЬ СРЕДЫ КАК ПРЕДПОСЫЛКА ДЛЯ УПРАВЛЕНИЯ СОСТОЯНИЕМ ПОПУЛЯЦИЙ 132  
A. P. Korzh, I. A. Akimov, T. V. Zahovalko
- ВПЛИВ ПРОБІОТИКА НА ПРОДУКТИВНІСТЬ КУРЧАТ-БРОЙЛЕРІВ 141  
Yu. M. Podolian
- ВИВЧЕННЯ ГОСТРОЇ ТОКСИЧНОСТІ ТА АНАЛГЕТИЧНОЇ АКТИВНОСТІ НОВИХ СПРОЦИКЛІЧНИХ ПОХІДНИХ ПІРАЗОЛІДИН-3,5-ДІОНУ 149  
V. M. Holovatiuk, O. A. Brazhko, V. I. Kashkovsky
- A NEW WEEVIL TRIBE, MEKORHAMPHINI TRIB. NOV. (COLEOPTERA, ITHYCERIDAE) WITH TWO NEW GENERA IN BURMESE AMBER 157

G. O. Poinar, A. E. Brown, A. A. Legalov CARPENTER-MOTHS OF TURKMENISTAN (LEPIDOPTERA: COSSIDAE) R. V. Yakovlev, T. J. Witt	165
ОЦІНКА ІСНУЮЧИХ ЗАГРОЗ ВІТРОПАРКІВ ПІВДНЯ УКРАЇНИ ДЛЯ МІГРАЦІЙНОЇ ОРНИТОФАУНИ	175
P. I. Gorlov, V. D. Siokhin, A. V. Matsyura ДОПОЛНЕНИЕ К ФЛОРЕ АЛТАЯ. II. A. A. Kechajkin, S. V. Smirnov, A. I. Shmakov, A. P. Shalimov, A. V. Vaganov, A. A. Batkin, M. V. Skaptsov, M. G. Kutsev, D. N. Shaulo, A. V. Matsyura	187
РАЗРАБОТКА ВЫСОКОАКТИВНОГО ШТАММА-ПРОДУЦЕНТА ВИРДЖИНИАМИЦИНА И ПОВЫШЕНИЕ ЕГО ПРОДУКТИВНОСТИ С ПОМОЩЬЮ СИНТЕТИЧЕСКИХ АБСОРБИРУЮЩИХ СМОЛ	195
V. A. Savushkin, V. V. Dzhavakhiya, E. V. Glagoleva, V. V. Savel'eva, E. D. Popova, A. I. Ovchinnikov, V. I. Glagolev, N. V. Novak, D. A. Durnikin МОРФОГЕНЕЗ ЯРОВОЙ ТВЕРДОЙ ПШЕНИЦЫ В КУЛЬТУРЕ ЗРЕЛЫХ ЗАРОДЫШЕЙ O. V. Bychkova, L. P. Khlebova, D. V. Ereschenko	209
РАЗРАБОТКА СУХОГО БАКТЕРИАЛЬНОГО ПРЕПАРАТА ДЛЯ СИЛОСОВАНИЯ С ОПТИМАЛЬНЫМ СООТНОШЕНИЕМ МОЛОЧНОКИСЛЫХ И ПРОПИОНОВОКИСЛЫХ КУЛЬТУР	219
M. A. Kartashov, T. M. Voinova, A. V. Sergeeva, N. V. Statsyuk, S. V. Rogovsky, Ya. O. Grebeneva, D. A. Durnikin ИССЛЕДОВАНИЕ РАЗМЕРА ГЕНОМА ПРЕДСТАВИТЕЛЕЙ РОДА POTENTILLA L. (ROSACEAE JUSS.)	229
A. A. Kechaykin, M. V. Skaptsov, S. V. Smirnov, M. G. Kutsev, A. I. Shmakov МАСШТАБНО-ЗАВИСИМЫЕ ЭФФЕКТЫ В СТРУКТУРЕ ЭКОЛОГИЧЕСКОЙ НИШИ ЛЕБЕДЯ-ШИПУНА CYGNUSOLOR (GMELIN, 1803) В ПЕРИОД ЗИМОВКИ В ПРЕДЕЛАХ ЗАЛИВА СИВАШ	234
A. Yu. Andrushenko, A. V. Zhukov ВОДОПРОНИКНІСТЬ ТЕХНОЗЕМІВ У МОДЕЛЯХ КОНСТРУКЦІЇ З РІЗНОЮ СТРАТИГРАФІЄЮ	250
K. P. Maslikova, I. V. Ladska, O. V. Zhukov ПРОДУКТИВНІСТЬ ТА ЗАБІЙНІ ЯКОСТІ ПЕРЕПЕЛІВ ПРИ РІЗНИХ РІВНЯХ ГУАНІДИНОЦТОВОЇ КИСЛОТИ В РАЦІОНАХ	266
M. Yu. Sychov, H. I. Pryumak WHAT IS LYCLENE DASARA (MOORE, 1859) (LEPIDOPTERA, EREBIDAE, ARCTIINAE)? Anton V. Volynkin, Karol Bucsek	275
CONTRIBUTION TO THE KNOWLEDGE OF THE LEPIDOPTERA IN THE NORTHERN CISOKHOTIA, RUSSIA	283
Matjaž Černila RESTORATION OF THE GENUS GURNA SWINHOE, 1892 (LEPIDOPTERA, EREBIDAE, ARCTIINAE)	290
Anton V. Volynkin THE DEVELOPMENT OF THE CONCEPT OF PLANT FUNCTIONAL TYPES WITH REGARD TO RARE SPECIES	295
A. A. Klimenko A REVIEW OF THE BARSINE FLAMMEALIS SPECIES-COMPLEX WITH DESCRIPTION OF A NEW SPECIES FROM HIMALAYA (LEPIDOPTERA, EREBIDAE, ARCTIINAE)	303
Anton V. Volynkin, Karel Černý	

ПШЕНИЦА МЯГКАЯ ОЗИМАЯ: ЭКОЛОГИЧЕСКАЯ ПЛАСТИЧНОСТЬ БИОЛОГИЧЕСКИХ И ТЕХНОЛОГИЧЕСКИХ МАРКЕРОВ	311
V. V. Moskalets, T. Z. Moskalets, S. P. Vasylykivskyi, I. V. Grynyk, A. G. Vovkohon, O. V. Lobova, O. A. Shevchuk, O. V. Knyazyuk	
POSSIBILITIES OF DE NOVO TRANSCRIPTOME SEQUENCING IN PYLOGENETIC RESEARCH ON AN EXAMPLE OF TARAXACUM OFFICINALE (ASTERACEAE)	319
M. G. Kutsev, M. V. Skaptsov, S. V. Smirnov, T. A. Sinitsyna, A. A. Kechaykin, M. S. Ivanova, A. I. Shmakov	
NEW DATA ON SOME OF THE RECENTLY DESCRIBED LITHOSIINI SPECIES (LEPIDOPTERA, EREBIDAE, ARCTIINAE)	324
Anton V. Volynkin, Vladimir V. Dubatolov, Karol Bucsek	
СПРАВНИТЕЛЬНОЕ ИССЛЕДОВАНИЕ МОРФОЛОГИИ СПОР РОДОВ ПОДСЕМЕЙСТВА CRYPTOGRAMMOIDEAE	333
A. V. Vaganov	
ГЕНОТОКСИЧЕСКИЕ ЭФФЕКТЫ ДОННЫХ ОТЛОЖЕНИЙ РЕКИ ЧУМЫШ В КЛЕТКАХ КОРНЕВОЙ МЕРИСТЕМЫ ALLIUM CEPA L.	347
L. P. Khlebova, N. N. Chernysheva, O. V. Bychkova, A. P. Kraynov	
REDESCRIPTION AND NEW DATA ON THE DISTRIBUTION OF A POORLY KNOWN PIRATE SPIDER ERMETUS INOPINABILIS (ARANEI: MIMETIDAE)	359
Yuri M. Marusik, Alexander A. Fomichev, Seppo Koponen	
ОЦЕНКА ЭФФЕКТИВНОСТИ ПРИМЕНЕНИЯ БИОАКУСТИЧЕСКОГО ОТПУГИВАТЕЛЯ ПТИЦ ДЛЯ УПРАВЛЕНИЯ ЧИСЛЕННОСТЬЮ ПТИЦ НА ТЕРРИТОРИИ ПОЛИГОНА ТВЕРДЫХ БЫТОВЫХ ОТХОДОВ ГОРОДА БАРНАУЛА	365
V. V. Shcherbinin, E. V. Ponkina, P. N. Ulanov, A. V. Matsyura	
АКТИВНІСТЬ ФЕРМЕНТІВ У ГЕПАТОПАНКРЕАСІ КОРОПА (CYPRINUS CARPIO L.) ЗА ДІЇ СУЛЬФАНІЛАМІДУ	377
Y. N. Kurbatova, V. M. Mykhalskaia, L. V. Malyuga, L. V. Gayova	
БІОХІМІЧНІ ТА МОРФОЛОГІЧНІ ПОКАЗНИКИ КРОВІ ЯПОНСЬКИХ ПЕРЕПЕЛІВ ЗА РІЗНИХ РІВНІВ КАЛЬЦІЮ ТА ФОСФОРУ У КОМБІКОРМАХ	383
L. M. Zlamanyuk	
ОСОБЛИВОСТІ ФОРМУВАННЯ ФІТОКЛІМАТУ ПІДКРОНОВОГО ПРОСТОРУ КУЩОВИХ ВИДІВ І ФОРМ ЯЛІВЦІВ В МЕЖАХ КОМПЛЕКСНОЇ ЗЕЛЕНОЇ ЗОНИ МІСТА ЛЬВОВА	390
T. I. Shuplat, V. V. Popovich	
МОРФОЛОГІЧНІ ОСОБЛИВОСТІ СЕЛЕЗІНКИ ТЕПЛОКРОВНИХ ТВАРИН	399
O. F. Dunaievskia	
ДИНАМІКА ПОПУЛЯЦІЇ ALLIUM OBLIQUUM L. В УМОВАХ ЗАХІДНОГО ПОДІЛЛЯ	407
N. V. Rubanovska, A. N. Solonenko, V. A. Solomakha	
ЦИАНОПРОКАРИОТЫ УРОЧИЩА ТУБАЛЬСКИЙ ЛИМАН	414
L. I. Arabadzhi, A. M. Solonenko, O. G. Bren, M. I. Holubev	
ANALYSIS OF THE ALIEN FLORA OF DNIPROPETROVSK PROVINCE	419
V. Baranoski, N. Khromykh, L. Karmyzova, I. Ivanko, Y. Lykholat	
АДВЕНТИВНЫЕ ВИДЫ ДРЕВЕСНЫХ РАСТЕНИЙ УРБАНОФЛОРЫ РОСТОВА-НА- ДОНУ	430
V. L. Kozlovsky, M. V. Kuropyatnikov, O. I. Fedorinova, M. M. Sereda, O. A. Kapralova, P. A. Dmitriev, T. V. Varduni	
NEW DATA ON THE CENTIPEDE (CHILOPODA) FAUNA OF EAST KAZAKHSTAN REGION	

Yu. V. Dyachkov, G. Sh. Farzaliyeva, A. A. Fomichev	438
СУДЕБНАЯ ЭНТОМОЛОГИЯ – ОБЛАСТЬ ЗНАНИЙ, НЕЗАСЛУЖЕННО ИГНОРИРУЮЩАЯСЯ В РОССИИ	443
R. V. Yakovlev, A. V. Matsyura, E. V. Guskova	
EGG SHAPE IN THE TAXONOMY AND PHYLOGENY OF BIRDS OF PREY	
I. S. Mytiai, V. I. Strigunov, A. V. Matsyura	447
BIRD SPECIES DATABASE OF EGG VOLUMES AND SURFACE AREAS	
I. S. Mytiai, A. V. Matsyura	455
MATHEMATICAL MODEL OF BIRD SPECIES IDENTIFYING: IMPLICATION OF RADAR DATA PROCESSING	463
V. V. Osadchyi, V. S. Ereemeev, A. V. Matsyura	
ВИКОРИСТАННЯ ОБЛПІХИ КРУШИНОВИДНОЇ ДЛЯ ФІТОРЕМЕДІАЦІЇ НАФТОЗАБРУДНЕНИХ ҐРУНТІВ	472
L. Z. Shevchyk, O. I. Romaniuk	
<b>Research Notes</b>	
SPECIES DIVERSITY AND DISTRIBUTION OF ENTOMOFAUNA IN THE SOUTHERN MIDDLE TAIGA OF THE MIDDLE OB	51
Z. I. Tyumaseva, E. V. Guskova	
SPECIES LIST OF PLUME MOTHS (LEPIDOPTERA, PTEROPHORIDAE) OF PARAGUAY WITH NEW RECORDS	88
P. Ustjuzhanin, V. Kovtunovich, A. Ustjuzhanina	
TO THE SYSTEMATICS AND NOMENCLATURE OF SOME PALAEARCTIC NYMPHALIDAE (LEPIDOPTERA)	118
S. K. Korb	
POLITZARIELLA FONTAINI – NEW COSSIDAE (LEPIDOPTERA) SPECIES FROM CONGO	154
R. V. Yakovlev, T. J. Witt	
NEW RECORDS OF DONACIA FABRICIUS, 1775 (COLEOPTERA: CHRYSOMELIDAE: DONACIINAE) FROM KRASNOYARSKII KRAI (RUSSIA)	
E. V. Guskova, E. N. Akulov	

L.I. Arabadzhi<sup>1</sup>, A.M. Solonenko<sup>1</sup>, O.G. Bren<sup>1,2</sup>, M.I. Holubev<sup>3</sup>  
**CYANOPROCARYOTA OF TUBALSKYI ESTUARY (AZOV SEA BASIN)**

<sup>1</sup>*Bogdan Chmelnytskyi Melitopol State Pedagogical University, Melitopol, Ukraine*

<sup>2</sup>*Pryazov National Natural Park, Melitopol, Ukraine*

<sup>3</sup>*National University of Life and Environmental Sciences of Ukraine, Kiev, Ukraine*

*E-mail: anatol8@ukr.net*

Information about species composition of Cyanoprocaryota representatives in the natural boundary of Tubalskyi Estuary within Pryazov National Natural Park is supplemented. Taxonomic structure of identified algae species is given, the data about occurrence of species in different habitats within national park is provided. We present a systematic structure and geographical affinity of discovered algae species. We registered 38 species of cyanoprocaryotic algae of orders: Chroococcales, Oscillatoriales and Nostocales with predominance of Oscillatoriales representatives in different habitats of Tubalskyi Estuary within the areas of Pryazov National Natural Park. The greatest number of algae species had genera *Lyngbya*, *Phormidium*, *Microcoleus* *Leptolyngbya* – 4 species per each genera. The majority of algae species were the cosmopolitans. *Microcoleus tenerrimus* was registered in all the studied plots of soil sampling and in the water reservoir of Tubalskyi Estuary. The most abundant and distributed species was *Lyngbya aestuarii*, which formed the macroscopic growths (cyanobacterial mats).

*Key words: Cyanoprocaryota, algae, Tubalskyi Estuary, Northern Pryazovye, national park.*

Л.І. Арабаджи<sup>1</sup>, А.М. Солоненко<sup>1</sup>, О.Г. Брен<sup>1,2</sup>, М.І. Голубев<sup>3</sup>  
**ЦИАНОПРОКАРИОТИ УРОЧИЩА ТУБАЛЬСЬКИЙ ЛИМАН**

<sup>1</sup>*Мелітопольський державний педагогічний університет імені Богдана Хмельницького*

<sup>2</sup>*Приазовський національний природний парк*

<sup>3</sup>*Національний університет біоресурсів і природокористування України*

*E-mail: anatol8@ukr.net*

Доповнені відомості видового складу представників відділу *Cyanoprocaryota* урочища Тубальський лиман в межах Приазовського національного природного парку. Наводиться систематична структура виявлених видів водоростей, надано дані щодо трапляння видів в різних біотопах урочища.

*Ключові слова: Cyanoprocaryota, водорості, Тубальський лиман, Північне Приазов'я, національний парк.*

Л.И. Арабаджи<sup>1</sup>, А.Н. Солоненко<sup>1</sup>, А.Г. Брен<sup>1,2</sup>, М.И. Голубев<sup>3</sup>  
**ЦИАНОПРОКАРИОТЫ УРОЧИЩА ТУБАЛЬСКИЙ ЛИМАН**

<sup>1</sup>*Мелитопольский государственный педагогический университет имени Богдана Хмельницкого*

<sup>2</sup>*Приазовский национальный природный парк*

<sup>3</sup>*Национальный университет биоресурсов и природопользования Украины*

*E-mail: anatol8@ukr.net*

Дополнены сведения видового состава представителей отдела *Cyanoprocaryota* урочища Тубальський лиман в пределах Приазовского национального природного парка. Приводится систематическая структура выявленных видов водорослей, предоставлены данные по встречаемости видов в различных биотопах урочища.

*Ключевые слова: Cyanoprocaryota, водоросли, Тубальский лиман, Северное Приазовье, национальный парк.*

**Citation:**

Arabadzhi, L.I., Solonenko, A.M., Bren O.G., Holubev, M.I. (2016). Cyanoprocaryota of Tubalskyi Estuary (Azov Sea basin). *Biological Bulletin of Bogdan Chmelnytskyi Melitopol State Pedagogical University*, 6 (3), 414–418.

**Поступило в редакцию / Submitted:** 18.11.2016

**Принято к публикации / Accepted:** 19.12.2016

**crossref** <http://dx.doi.org/10.15421/2016112>

© Arabadzhi, Solonenko, Bren, Holubev, 2016

Users are permitted to copy, use, distribute, transmit, and display the work publicly and to make and distribute derivative works, in any digital medium for any responsible purpose, subject to proper attribution of authorship.



This work is licensed under a Creative Commons Attribution 3.0. License

North-Western territories of the Azov Sea shore have unique landscapes and high biodiversity is extremely valuable in the economical aspect and have great recreational potential. In 2010 by decree of President of Ukraine the environmental institution – Pryazov National Natural Park (hereinafter - PNPP) was established for the rational use and conservation of natural resources in this area (Chronicles..., 2012). One of the main objectives of environmental institutions (including PNPP) is to carry out the monitoring researches of natural ecosystems. The cyanoprocaryotic algae could be used as the convenient and reliable bioindicators of processes in ecosystems. The results of researches of algae from the North-Western territories of the Azov Sea shore were presented in several articles (Solonenko et al., 2006, 2008), but cyanoprocaryotic algae within the natural boundary of Tubalskyi Estuary are studied unevenly and should be updated by additional data.

Natural boundary of Tubalskyi Estuary is situated near the Village Prymorskyi Posad. Its' territories have a flat landscape and formed by the mouth of River Domuzla (Fig. 1). The maximum length of the tract - 4.15 km, maximum width - 2 km, altitude of slopes of the right bank - 9 m. The coordinates of extreme points: North - 46°37'5.93"N 35°42'11.27"E; South - 46°34'51.04"N 35°43'4.87"E; West - 46°36'18.83"N 35°40'49.32"E; East - 46°35'46.97"N 35°43'58.63"E. Water regime depends mainly on precipitation as well as river water, sea water infiltration, periodic flooding through the sand bar. The maximum length of water - 300 m, maximum width - 100 m, maximum depth - 25 cm. The water surface area greatly varies and depends on the season. The salinity of the water is 55-112 ‰; pH 6.1-6.4. Tubalskyi Estuary is included in the list of protected areas with a status of the wetland of regional importance.

The higher vegetation is presented here by plant communities with dominance of *Phragmites australis* (Cav.) Trin. ex Steud. and halophilic communities with dominance of *Salicornia europea* L., *Halocnemum strobilaceum* M. Bieb., *Salsola soda* L., *Suaeda altissima* (L.) Pall. There is a pond in the southern part of the natural boundary, and in the warmer months the *Potamogeton pectinatus* (L.) Börner develops powerful growths over there.



Figure 1. Map of sampling points

## MATERIALS AND METHODS

The soil samples were carried out in the 2014–2015 on the four fixed sampling points in the natural boundary of Tubalskyi Estuary within the territory of PNNP (Pryazov district, Zaporizhzhya region), namely: steppe slopes, sandy soils, saline soils and in the reservoir. Samples were collected in spring, summer and autumn by the conventional phycology methods (Hollerbach, 1969; Topachevskiy, 1984). Totally, we sampled and processed 18 terrestrial samples and 9 water samples. The subsequent identification of algae was performed at the Department of Botany and Landscape Architecture in the laboratory of algal and ecological studies of the terrestrial and aquatic ecosystems in the Bogdan Chmelnytskyi Melitopol State Pedagogical University.

Collected material was processed using cultural methods: 1) soil culture with glasses of growth; 2) agaric Bold's nutrient media with normal and tripled quantity of nitrogen (1N BBM and 3N BBM, respectively) (Arce, 1958). Researches were held on by the light binocular microscope "MICROmed XS-5520" using 40 $\times$  and 100 $\times$  lenses. Identification was carried out by the identification guide of algae (Komárek, Anagnostidis, 1998; 2005). Analysis of the revealed algae species in relation to some geographical confinement was carried out according to Barinova (2006). Systematic list of the identified algae species was structured by system adopted in the monograph of I. Kostikov (2001) with additions and clarifications according to a check-list by Wasser (2000) and Tsarenko (2006).

## RESULTS AND DISCUSSION

We identified 38 species of algae in the natural boundary of Tubalskyi Estuary. These species belongs to 3 orders, 12 families and 19 genera (Table. 1).

Table 1. Systematic structure of cyanoprocarotic algae, Tubalskyi Estuary

№	Taxon
Phylum	<b>Cyanoprocarota</b>
Class	Cyanophyceae Sachs 1874
Order	Chroococcales Cavalier-Smith, 2002
Family	<i>Merismopediaceae</i> Elenkin 1933
Genus	<i>Synechocystis</i> Sauvageau 1892
1.	<i>Synechocystis salina</i> Wislouch 1924
2.	<i>Synechocystis crassa</i> Woronichin 1929
Genus	<i>Merismopedia</i> Meyen 1839
3.	<i>Merismopedia elegans</i> A. Braun in Kützing 1849
4.	<i>Merismopedia glauca</i> Kützing 1845
5.	<i>Merismopedia punctata</i> Meyen 1839
Family	<i>Microcystaceae</i> Elenkin 1933
Genus	<i>Microcystis</i> Kützing 1907
6.	<i>Microcystis pulverea</i> (Wood) Forti emend Elenkin 1933
Genus	<i>Chondrocystis</i> Lemmermann 1899
7.	<i>Chondrocystis sarcinoides</i> Komárek et Anagnostidis 1995
Family	<i>Chroococcaceae</i> Nägeli 1849
Genus	<i>Pseudocapsa</i> Ercegović 1925
8.	<i>Pseudocapsa sphaerica</i> Kováčik 1988
Family	<i>Entophysalidaceae</i> Geitler 1925
Genus	<i>Johannesbaptistia</i> De Toni 1934
9.	<i>Johannesbaptistia pellucida</i> Taylor et Drouet 1938
Family	Hyellaceae Borzi 1914
Genus	<i>Hyella</i> Bornet et Flahault 1886
10.	<i>Hyella caespitosa</i> Bornet et Flahault 1888
Order	Oscillatoriales Elenkin 1934
Family	<i>Oscillatoriaceae</i> Engler 1898
Genus	<i>Lynghya</i> Agardh 1892
11.	<i>Lynghya aestuarii</i> Liebman ex Gomont 1892
12.	<i>Lynghya semiplena</i> J. Agardh ex Gomont 1892
13.	<i>Lynghya salina</i> Kutzing ex Starmach 1966



№	Taxon
14.	<i>Lyngbya major</i> Meneghini 1892
Genus	<i>Oscillatoria</i> Vaucher 1892
15.	<i>Oscillatoria tenuis</i> Agardh ex Gomont 1892
16.	<i>Oscillatoria geminata</i> Meneghini 1892
Family	<i>Phormidiaceae</i> Anagnostidis et Komárek 1988
Genus	<i>Phormidium</i> Kützing 1892
17.	<i>Phormidium uncinatum</i> (Agardh) Gomont 1890
18.	<i>Phormidium autumnale</i> Gomont 1892
19.	<i>Phormidium paulsenianum</i> Boye-Petersen 1930
20.	<i>Phormidium takyricum</i> Novitschkova 1960
Genus	<i>Pseudophormidium</i> Anagnostidis et Komárek 1988
21.	<i>Pseudophormidium edaphicum</i> (Elenkin) Anagnostidis et Komárek 1988
Genus	<i>Microcoleus</i> Desmazières 1892
22.	<i>Microcoleus vaginatus</i> Gomont 1892
23.	<i>Microcoleus paludosus</i> (Kützing) Gomont 1892
24.	<i>Microcoleus tenerimus</i> Gomont 1892
25.	<i>Microcoleus chthonoplastes</i> Thuret ex Gomont 1892
Genus	<i>Spirulina</i> Turpin ex Gomont 1892
26.	<i>Spirulina subsalsa</i> Oersted ex Gomont 1892
Family	<i>Plectonemataceae</i> Elenkin 1934
Genus	<i>Plectonema</i> Thuret 1892
27.	<i>Plectonema notatum</i> (Schmidle) Anagnostidis & Komárek 1988
Genus	<i>Symploca</i> Kützing 1892
28.	<i>Symploca elegans</i> Kützing 1892
Family	<i>Schizotrichaceae</i> Elenkin 1934
Genus	<i>Schizothrix</i> Kützing 1892
29.	<i>Schizothrix coriacea</i> Kützing ex Gomont 1892
30.	<i>Schizothrix friesii</i> (Agardh) Gomont 1892
31.	<i>Schizothrix lardacea</i> (Cesati) Gomont 1892
Family	<i>Pseudanabaenaceae</i> Anagnostidis et Komárek 1988
Genus	<i>Leptolyngbya</i> Anagnostidis et Komárek 1988
32.	<i>Leptolyngbya valderiana</i> (Gomont) Anagnostidis et Komárek 1988
33.	<i>Leptolyngbya halophila</i> (Hansgirg ex Gomont) Anagnostidis et Komárek 1988
34.	<i>Leptolyngbya boryana</i> (Gomont) Anagnostidis et Komárek 1988
35.	<i>Leptolyngbya notata</i> (Schmidle) Anagnostidis et Komárek 1988
Order	Nostocales Cavalier-Smith 2002
Family	<i>Nostocaceae</i> Eichler 1886
Genus	<i>Nostoc</i> Vaucher 1886
36.	<i>Nostoc punctiforme</i> (Kützing) Hariot 1891
37.	<i>Trichromus ellipsosporus</i> (Fritsch) Komárek et Anagnostidis 1989
Family	<i>Rivulariaceae</i> Kützing 1843
Genus	<i>Calothrix</i> Agardh 1886
38.	<i>Calothrix parietina</i> Thuret 1886

The greatest species diversity had order Oscillatoriales (21 species) followed by *Phormidiaceae* – 10 species, *Lyngbya*, *Phormidium*, *Microcoleus* *Leptolyngbya* – 4 species. Such genera as *Merismopedia*, *Schizothrix* and *Synechocystis*. *Oscillatoria* were represented by 3 and 2 species respectively; other genera – by 1 species. According to literature data 19 identified algae species could be related to cosmopolitan (16) and Holarctic species (3). Among all studied habitats the largest number of cyanoprocaryotic algae (15 species) were found on the steppe slopes: *Leptolyngbya boryana*, *Lyngbya semiplena*, *Microcoleus paludosus*, *Microcoleus tenerimus*, *Microcoleus vaginatus*, *Nostoc punctiforme*, *Oscillatoria*

*tenuis*, *Phormidium uncinatum*, *Pseudophormidium edaphicum*, *Schizothrix coriacea*, *Schizothrix friesii*, *Schizothrix lardacea*, *Symploca elegans*, *Plectonema notatum*, *Trichromus ellipsosporus*. Algae of the steppe slopes formed local greening-like macroscopic growths on the ground surface, which were observed in the lower parts of the slopes in the places of moisture accumulation. Also the great species variety was registered in saline soils (solonchak) and in the reservoir – 11 species per each habitat. In the salt marshes we founded such species as: *Leptolyngbya halophila*, *Leptolyngbya valderiana*, *Lyngbya major*, *Lyngbya aestuarii*, *Lyngbya salina*, *Lyngbya semiplena*, *Microcoleus chthonoplastes*, *Microcoleus tenerimus*, *Phormidium paulsenianum*, *Phormidium takyricum*, *Schizothrix coriacea*. *Lyngbya aestuarii* was the only species which formed the dark-green solid slime of macroscopic growths.

We identified *Chondrocystis sarcinoides*, *Hyella caespitosa*, *Johannesbaptistia pellucida*, *Merismopedia punctata*, *Microcystis pulvereae*, *Microcoleus chthonoplastes*, *Microcoleus tenerimus*, *Pseudocapsa sphaerica*, *Synechocystis crassa*, *Synechocystis salina*, *Spirulina subsalsa* for the area of Tubalskyi Estuary whereas the characteristic algal water blooms were not observed in warm periods. The limited number of blue-green algae species were sampled in sandy soils: *Calothrix parietina*, *Leptolyngbya notata*, *Lyngbya semiplena*, *Merismopedia elegans*, *Merismopedia glauca*, *Oscillatoria tenuis*, *Oscillatoria geminata*, *Phormidium autumnale* where the algae did not form the massive growths during the research period. The most abundant and frequent species in all the habitats were *Microcoleus tenerimus*, *Microcoleus chthonoplastes*, *Lyngbya semiplena*, *Schizothrix coriacea*, while *Microcoleus tenerimus* was registered in all studied habitats of terrestrial and water zone of Tubalskyi Estuary area.

### CONCLUSIONS

We registered 38 species of cyanoprokaryotic algae of orders: Chroococcales, Oscillatoriales and Nostocales with predominance of Oscillatoriales representatives in different habitats of Tubalskyi Estuary within the areas of Pryazov National Natural Park. The greatest number of algae species had genera *Lyngbya*, *Phormidium*, *Microcoleus* *Leptolyngbya* – 4 species per each genera. The majority of algae species were the cosmopolitans. *Microcoleus tenerimus* was registered in all the studied plots of soil sampling and in the water reservoir of Tubalskyi Estuary. The most abundant and distributed species was *Lyngbya aestuarii*, which formed the macroscopic growths (cyanobacterial mats).

### REFERENCES

- Arce, G., Bold, H.C. (1958). Some Chlorophyceae from Cuban Soils. Amer. Bot. Journ, 45, 492-503.
- Baraboha, N.M., Baraboha, A.P., Bren, O.G. (2012). Chronicles of Pryazov National Natural Park (2011). Pryazov National Park. Melitopol (in Russian).
- Barinova, S.S., Medvedeva, L.A., Anisimova, O.V. (2006). Biodiversity of Environmental Indicator Algae. Tel Aviv: Pilies Studio.
- Hollerbach, M.M., Shtina, E.S. (1969). Soil algae. Leningrad: Nauka (in Russian).
- Komarek, J., Anagnostidis, K. (1999). Cyanoprokaryota. 1 Teil: Chroococcales. In: Ettl H., Gerloff J., Heynig H., Mollenhauer D. (eds), Süßwasserflora von Mitteleuropa, Gustav Fischer Verlag.
- Komárek, J., Anagnostidis, K. (2005). Cyanoprokaryota I. Oscillatoriales. Süßwasserflora von Mitteleuropa. – Jena-Stuttgart-Lübeck-Ulm: Gustav Fischer.
- Kostikov, I.Yu., Romanenko, P.A., Demchenko, E.M. (2001). Soil algae of Ukraine (history and methods, system synopsis of flora). Kiev: Fitosotsiotsentr (in Russian).
- Solonenko, A.N., Yarovoy, S.A., Podorozhnyi, S.N., Raznopolov, O.N. (2006). Algae from the solonchak of the Stepanovska and Fedotova Spits in the North-West coast of the Azov Sea. Soil Science, 7(3-4), 123-127. (in Russian).
- Solonenko, A.N., Yarovoy, S.A., Yarovaya, T.A. (2008). Algae from the solonchak of the mouth parts of river Korsak in the natural boundary Tubalskyi estuary. Bulletin of the State Nikitsky Botanical Garden, 96, 26-29 (in Russian).
- Topachevskiy, A.V., Masuk, N.P. (1984). Freshwater algae of Ukrainian SSR. Kiev: Vyshaya Shkola (in Russian).
- Tsarenko, P.M., Wasser, S.P., Nevo, E. (2006). Algae of Ukraine. Diversity, Nomenclature, Taxonomy, Ecology and Geography. Ruggell: A.R.A. Gantner Verlag K.G.
- Wasser, S.P., Tsarenko, P.M. (2000). Algae diversity of Ukraine. Phycology, 10(4), 1-309.
- Yarovaya, T.A., Yarovoy, S.O., Bren, O.G. (2012). Algae of the Pryazov National Natural Park. Proceed. Int. Conf. The role of environmental institutions in preserving biodiversity, ethnic and cultural heritage and sustainable development areas. CBS: PNP Huzulschyna (in Ukrainian).
- Yarovoy, S.A., Yarovaya, T.A., Solonenko, A.N. (2008a). For the study of the solonchak algae on the Berdyansk Spit near the lake Krasnoye. Ecology and Noospherology, 19(1-2), 160-162 (in Russian).
- Yarovoy, S.A., Solonenko, A.N., Yarovaya, T.A. (2008b). Macroscopic growths of algae on the sea shore solonchaks of the Azov Sea. Proceed. Int. Sc. Conf. and Marine Biology School "Modern problems of phycology". Rostov-on-Don (in Russian).
- Yarovoy, S.A. (2012). Macroscopic growths of algae on the solonchaks of the Priazov National Natural Park (Ukraine). Proceed. IV Int. Conf. Actual problems of modern phycology. Kiev (in Russian).
- Yarovoy, S.A., Solonenko, A.N., Olejnik, T.A. (2007). Soil algae of the sea shore solonchak of the Berdyansk Spit near the lake Krasnoye. Proceed. Int. Conf. Biology of the XXI century: theory, practice and teaching. Cherkasy–Kaniv (in Russian).
- Yarovy, S.O., Yarovaya, T.A., Bren, O.G. (2013). Algological study of the Pryazov National Natural Park wetlands. In: Ecology of wetlands and peatlands. Kyiv: Dia (in Ukrainian).